CORRECTION Open Access



## Correction to: Fetal growth is associated with CpG methylation in the P2 promoter of the *IGF1* gene

Catherine Le Stunff<sup>1\*</sup>, Anne-Laure Castell<sup>2</sup>, Nicolas Todd<sup>1</sup>, Clémence Mille<sup>1</sup>, Marie-Pierre Belot<sup>1</sup>, Nathalie Frament<sup>1</sup>, Sylvie Brailly-Tabard<sup>3</sup>, Alexandra Benachi<sup>4</sup>, Delphine Fradin<sup>5</sup> and Pierre Bougnères<sup>1</sup>

## Correction

After publication of the original article [1], it came to the publishers' attention that the below author's corrections provided at the proofing stage had been misinterpreted.

- 1) The title should have read "Fetal growth is associated with CpG methylation in the P2 promoter of the *IGF1* gene", and not "Fetal growth is associated with the CpG methylation of the P2 promoter of the *IGF1* gene".
- 2) The first sentence of Fig. 1 caption should have read "CpG-137 methylation correlates negatively with **birth length**", and not "CpG-137 methylation correlates negatively with **height**".
- 3) Additional file 3 legend should have read "Correlation matrix of methylation values (%) at the **CpG** located in the P1 and P2 promoters of the IGF1 gene in newborns patients. Pearson correlation coefficient is indicated in bold, and P value below.", and not "Correlation matrix of methylation values (%) at the **CG** located in the P1 and P2 promoters of the IGF1 gene in newborns patients. Pearson correlation coefficient is indicated in bold, and P value below."
- 4) Additional file 4 legend should have read "Relationship between promoter **CpG** methylation and genotypes. (A) Methylation at CpGs-137 of the IGF1 P2 promoter is independent from the rs35767 genotypes. (B) Methylation at CpGs-206 and CpG-180 in insulin promoter is closely dependent on rs689 alleles.", and not "Relationship between promoter **CG** methylation and genotypes. (A) Methylation at CpGs-137 of the IGF1 P2 promoter

is independent from the rs35767 genotypes. (B) Methylation at CpGs-206 and CpG-180 in insulin promoter is closely dependent on rs689 alleles."

The original article has been updated to reflect these corrections.

## **Author details**

<sup>1</sup>Institut National de la Santé et de la Recherche Médicale U1169, Bicêtre Hospital, Paris Sud University, Le Kremlin-Bicêtre, France. <sup>2</sup>Service de Médecine des Adolescents, Bicêtre Hospital, Paris Sud University, Le Kremlin-Bicêtre, France. <sup>3</sup>Service de BiologieMoléculaire et Hormonologie, Bicêtre Hospital, Paris Sud University, Le Kremlin-Bicêtre, France. <sup>4</sup>Service de Gynécologie-Obstétrique, Antoine Béclère Hospital, Paris Sud University, Clamart, France. <sup>5</sup>CRCINA, INSERM U1232, Université de Nantes, Nantes, France.

Received: 18 May 2018 Accepted: 18 May 2018 Published online: 04 June 2018

## Reference

 Le Stunff C, Castell A, Todd N, Mille C, Belot M, Frament N, Brailly-Tabard S, Benachi A, Fradin D. Fetal growth is associated with CpG methylation in the P2 promoter of the IGF1 gene. Clin Epigenetics. 2018;10:57. https://doi.org/ 10.1186/s13148-018-0489-9.

<sup>1</sup>Institut National de la Santé et de la Recherche Médicale U1169, Bicêtre Hospital, Paris Sud University, Le Kremlin-Bicêtre, France



<sup>\*</sup> Correspondence: catherine.le-stunff@inserm.fr