

Treatment of poor-risk myelodysplastic syndromes and acute myeloid leukemia with a combination of 5-azacytidine and valproic acid

Andrea Kuendgen · Gesine Bug · Oliver G. Ottmann · Detlef Haase · Julie Schanz · Barbara Hildebrandt · Kathrin Nachtkamp · Judith Neukirchen · Ariane Dienst · Rainer Haas · Ulrich Germing · Norbert Gattermann

Received: 1 December 2010 / Accepted: 20 March 2011 / Published online: 8 April 2011
© Springer-Verlag 2011

Abstract 5-azacytidine (AZA) has become standard treatment for patients with higher-risk myelodysplastic syndrome (MDS). Response rate is about 50% and response duration is limited. Histone deacetylase (HDAC) inhibitors are attractive partners for epigenetic combination therapy. We treated 24 patients with AZA (100 mg/m², 5 days) plus valproate (VPA; continuous dosing, trough serum level 80–110 µg/ml). According to WHO classification, 5 patients had MDS, 2 had MDS/MPD, and 17 had acute myeloid leukemia (AML). Seven patients (29%) had previously received intensive chemotherapy, and five had previous HDAC inhibitor treatment. The overall response rate was 37% in the entire cohort but significantly higher (57%) in previously untreated patients, especially those with MDS (64%). Seven (29%) patients achieved CR (29%) and two PR (8%), respectively. Hematological CR was accompanied by complete cytogenetic remission according to conventional cytogenetics

in all evaluable cases. Some patients also showed complete remission according to FISH on bone marrow mononuclear cells and CD34⁺ peripheral blood cells, as well as by follow-up of somatic mitochondrial DNA mutations. Four additional patients achieved at least marrow remissions. Factors influencing response were AML (vs. MDS), marrow blast count, pretreatment, transfusion dependency, concomitant medication with hydroxyurea, and valproic acid (VPA) serum level. This trial is the first to assess the combination of AZA plus VPA without additional ATRA. A comparatively good CR rate, relatively short time to response, and the influence of VPA serum levels on response suggest that VPA provided substantial additional benefit. However, the importance of HDAC inhibitors in epigenetic combination therapy can only be proven by randomized trials.

A. Kuendgen (✉) · K. Nachtkamp · J. Neukirchen · A. Dienst · R. Haas · U. Germing · N. Gattermann
Department of Hematology, Oncology and Clinical Immunology,
Heinrich-Heine-University,
Moorenstr. 5,
40225 Düsseldorf, Germany
e-mail: kuendgen@med.uni-duesseldorf.de

G. Bug · O. G. Ottmann
Department of Hematology and Oncology,
Johann Wolfgang Goethe University,
Frankfurt/Main, Germany

D. Haase · J. Schanz
Department of Hematology/Oncology,
Georg-August-University Göttingen,
Göttingen, Germany

B. Hildebrandt
Institute of Human Genetics, Heinrich-Heine-University,
Düsseldorf, Germany

Introduction

In recent years, epigenetic therapy has become a treatment option for patients with higher-risk myelodysplastic syndrome (MDS) who are not considered candidates for intensive induction chemotherapy or allogeneic stem cell transplantation (SCT). The demethylating agent 5-azacytidine (AZA) can achieve substantial survival benefit for patients with higher-risk MDS and patients with acute myeloid leukemia (AML) who have a bone marrow blast count of 20–30% (RAEB-T according to the FAB classification) (Fenaux et al. 2009). Although complete response (CR) rates are not higher than 10–20% (Fenaux et al. 2009; Silverman et al. 1994; Silverman et al. 2002, and Silverman et al. 2006), almost half of the patients with intermediate-II or high-risk disease according to IPSS (Greenberg et al. 1997) show hematological improvement. Responses are usually seen

only after several treatment cycles. Lengthy time to response is problematic for patients with an aggressive course of disease, particularly patients with AML. Results from phase II trials with azacitidine or decitabine suggest that only about one third of such patients respond (Lubbert et al. 2008; Maurillo et al. 2008). To further improve remission rates, time to response and response duration, combinations of AZA with other agents are being evaluated.

Since epigenetic treatment aims at reversing pathological gene silencing, and DNA methylation cooperates with histone modification to control gene expression, it appears logical to combine AZA with inhibitors of histone deacetylases. Preclinical studies suggest that pharmacologic targeting of both, DNA methyltransferases (DNMT) and histone deacetylases (HDAC), may result in synergistic anticancer activity (Bhalla 2005; Yang et al. 2005).

In 2001, two independent groups showed that the antiepileptic drug valproic acid (VPA) also has HDAC inhibitory activity and induces differentiation of malignant myeloid cells, an ability that is enhanced by all-trans retinoic acid (ATRA) (Göttlicher et al. 2001; Phiel et al. 2001). Stimulated by these findings, we studied the clinical effect of VPA at serum concentrations of 50–100 µg/ml in 23 patients with AML or MDS as monotherapy or in combination with (ATRA) (Kuendgen et al. 2004). The pilot study yielded an overall response rate of 35%. Interestingly, response rate was 44% for patients receiving VPA monotherapy, while none of five patients receiving VPA+ATRA from the start responded. Responses were more frequent in lower-risk MDS, but some patients with higher-risk MDS showed a decrease of their elevated blast count. Follow-up of 122 patients confirmed the higher response rates achieved in low-risk MDS. Only few patients with high-risk MDS benefited from VPA monotherapy or VPA+ATRA. Based on our experience with VPA (Kuendgen et al. 2004; Kuendgen et al. 2006, and Kuendgen and Gattermann 2007) and AZA (Fenaux et al. 2009), we embarked on evaluating the combination of the two drugs in patients with MDS and AML.

Patients and methods

Study design

Primary endpoint of the study was the feasibility and safety of a combination treatment with AZA plus VPA. Secondary endpoints were overall and progression-free survival, as well as hematological response rate according to revised International Working Group (IWG) criteria (Cheson et al. 2006). Study treatment was initiated with AZA 100 mg/m²/day for 5 days every 28 days administered subcutaneously. We chose to investigate a 5-day schedule which is easier to

apply than the approved 7-day schedule (75 mg/m²/day for days 1–7) while providing almost the same cumulative dose per cycle. Treatment with VPA was started on day 4. The dosage of daily oral VPA was adjusted to achieve trough serum concentrations between 80 and 110 µg/ml, i.e., in the upper therapeutic range for antiepileptic treatment. Serum VPA levels were measured with a commercially available fluorescence polarization immunoassay (Abbott, Wiesbaden, Germany). Response evaluation was performed by bone marrow cytology, conventional cytogenetics, FISH on bone marrow mononuclear cells (BM-FISH) and CD34⁺ cells in the peripheral blood (PB-CD34-FISH) (Bräulke et al. 2010), as well as analysis of somatic mitochondrial DNA mutations (Wulfert et al. 2008). Treatment was continued as long as neither significant side effects nor disease progression occurred. Evaluation of treatment response included all patients who had received at least one complete cycle of study medication and was scheduled at least 1 year after inclusion of the last patient. Two centers participated in the trial, namely the University Hospital of Heinrich-Heine-University, Düsseldorf and the University Hospital of Johann Wolfgang Goethe-University, Frankfurt/Main.

Inclusion criteria

Patients included in the study had either

- primary MDS with poor-risk features: bone marrow blast count $\geq 10\%$ and/or poor-risk karyotype according to IPSS (Greenberg et al. 1997) or
- therapy-related, secondary MDS or AML, or
- AML according to WHO criteria (Harris et al. 1999) or
- chronic myelomonocytic leukemia (CMML-II).

Inclusion criterion for all patients was relapse after, ineligibility for, or refusal of intensive chemotherapy or allogeneic SCT. Ineligibility for intensive treatment was determined by the treating physician after thorough discussion with the patient. Further inclusion criteria included age ≥ 18 years, Karnofsky index $>50\%$, negative pregnancy test, and written informed consent according to the Declaration of Helsinki. Patients were excluded if they were eligible for intensive chemotherapy or allogeneic SCT, had impaired liver function (bilirubin, ALT, and/or AST $>2\times$ upper limit of normal) and/or reduced renal function (creatinine $>2.5\times$ upper limit of normal), previous liver failure, liver or pancreatic disease, or a family history of liver failure, especially after treatment with VPA, porphyria, or any kind of coagulopathy.

Patient characteristics

Between March and July 2007, 25 patients were enrolled, of whom 24 received at least one complete cycle of therapy

and where thus evaluable for response. One patient could not receive a full cycle of vidaza and did not receive VPA due to very rapid disease progression. Median age was 73 years (59–87 years). According to FAB classification (Bennett et al. 1982), 13 patients had MDS and 11 patients suffered from AML. According to WHO (Harris et al. 1999), only 4 patients had MDS, 2 MDS/MPD, and 18 AML (6 de novo, 7 sAML/MDS, 5 AML/MDS therapy related). Only nine patients (37%) had low-risk cytogenetics according to IPSS (Greenberg et al. 1997), all of whom showed a normal karyotype. Four patients (17%) had intermediate-risk and 11 (46%) had high-risk cytogenetics, including seven patients with aberrations of chromosome 7 (2 as single aberration, 2 with one additional abnormality, and 3 as part of a complex karyotype). According to the Grimwade definition for AML (Grimwade et al. 1998), none of our patients had favorable cytogenetics, 12 (50%) had intermediate-risk karyotypes and 10 patients (50%) had adverse cytogenetics. Seventeen patients (71%) were considered ineligible for intensive chemotherapy, because of old age (>75 years) ($n=5$), concomitant disease ($n=6$), or adverse cytogenetics in patients >60 years of age ($n=8$). Seven patients (29%) had previously received one ($n=3$) or more ($n=4$) cycles of chemotherapy and had relapsed ($n=5$) or were refractory ($n=2$). Five patients had previously received treatment with an HDAC inhibitor (LBH589, $n=3$; VPA, $n=2$). Patient characteristics are summarized in Table 1.

Statistical analysis

The product-limit method was used to estimate the probability of survival. Survival was measured from start of treatment to time of death. Survival curves were compared using the log-rank test. Clinical and hematological parameters of patients at the time of treatment were compared using the chi-square test with Yates correction and the Wilcoxon rank-sum test. Parameters that were correlated with response were analyzed using logistic regression and stepwise multivariate regression method of Cox.

Results

Hematological response

Of the 24 patients evaluable for response, 7 achieved cytological CR (29%) and two had PR (8%), giving an overall response rate of 37%. All these patients showed hematologic improvement. One CR was a CRi (with incomplete recovery of platelets to less than 100,000/ μ l), and in five cases dysplastic features persisted (CRdys). In

Table 1 Patient characteristics

Characteristic	No.
Median age/years (range)	73 (59–87)
Gender ($n=$)	
Male/female	19/5
Diagnosis (FAB) ($n=$)	
RAEB	8
RAEB-T	3
CMML	2
AML	11
Diagnosis (WHO) ($n=$)	
RAEB II	4
CMML II	2
AML, de novo	6
sAML/MDS	7
AML/MDS, therapy-related	5
IPSS-risk group ($n=$)	
Intermediate II	5
High	9
Karyotype-risk (IPSS) ($n=$)	
Good	9
Intermediate	4
Poor	11
Karyotype risk (AML ¹) ($n=$)	
Good	0
Intermediate	12
Poor	12
Karyotype ($n=$)	
Normal	7
Abnormal	17
Median marrow blast count (%)	18 (1–80)
Pre-treatment ($n=$)	
Intensive chemotherapy	1
Hydroxyurea	3
LBH-589	3
VPA (+/- ATRA)	2
Growth factors	2
None	2
Transfusion dependency	
Yes/no	19/5
Median WBC/ μ l	2,245 (400–32,000)
WBC>10,000 ($n=$)	5
WBC<1,800	20
WBC<500	7
Median platelet count/ μ l	57,000 (5,000–252,000)
Platelets<100,000 ($n=$)	17
Platelets<30,000	9

FAB French–American–British classification, *WHO* World Health organization classification, *IPSS* International Prognostic Scoring System, *AML* acute myeloid leukemia, *VPA* valproic acid, *ATRA* all-trans retinoic acid, *WBC* white blood count. 1 Grimwade 1998

addition to these nine patients with CR/PR, four patients showed decreasing marrow blasts (three complete marrow responses and one reduction from 34 to 8%), though without hematologic improvement. These cases were not considered as responders.

Interestingly, patient 14 with sAML/MDS, whose marrow blasts decreased from 22 to 2% and who developed Coombs-negative hemolysis, responded well to erythropoietin which was started after the official end of the study (cycle 24). Despite a relatively high endogenous Epo level of 584 mU/l, the patient became transfusion independent (maximum hemoglobin 13.6 g/dl). After cycle 24, the blast count increased to 22% and then 50%. Surprisingly, peripheral blood counts remained stable and the patient was transfusion-independent for another 8 months despite full-blown relapse.

Cytogenetic response

In the six evaluable patients, hematological CR or PR was always accompanied by complete remission as assessed by conventional cytogenetics (CgCR). The cytogenetic diagnosis of the responders (+21; complex including monosomy 7; +8; del(7q); -7 plus dup(1q); del(20q)) included three poor-risk karyotypes.

Cytogenetic responses were also analyzed by FISH on bone marrow mononuclear cells (BM-FISH) and CD34⁺ cells in the peripheral blood (PB-CD34-FISH). Only the patient with CRi did not have BM-FISH. In the other five, complete remission was confirmed in four, while in one patient, 6% cells with monosomy 7 persisted among BM-MNC. However, in this patient, no aberrant cells were detectable by PB-CD34-FISH. Among the six patients with CgCR by conventional cytogenetics, four also showed CgCR on analysis of PB-CD34-FISH, while two achieved only CgPR (91 → 5 → 29%, and 79 → 1%).

Patient no. 6 showed an interesting development. Within 2 cycles, he achieved complete cytomorphic remission. After 5 cycles, he also showed CgCR on conventional cytogenetics and BM-FISH. At that time, 32% of his PB-CD34⁺ cells still showed del(7q) (from an initial 91%). After 9 cycles, the proportion had fallen to 5%. After the first 2 cycles of treatment, an additional clone characterized by loss of the Y chromosome was detected in the marrow, which had not been described prior to study treatment and increased to 53% of metaphases after 5 cycles. Thereafter, that clone decreased slowly to 0%, in parallel to an increase of del(7q) cells. A third clone was observed after 13 cycles exhibiting complete monosomy 7. Cytologically, the patient showed only a borderline blast count of 5% after 17 cycles. The astonishing observation in this case is that peripheral blood counts remained completely stable, and that the patient had never required transfusions for now 43 treatment cycles

despite recurrence of del(7q) and karyotype evolution with development of a second clone with monosomy 7.

Of the patients who achieved hematologic remission, only three had somatic mitochondrial DNA mutations suitable for follow-up. Two showed a CR (70 → 0 → 50%, and 67 → 0%) and one had a PR (50 → 5%). Characteristics of responders and patients with marrow remissions are shown in Table 2. Tables 3 and 4 present an overview of treatment details and responses.

Median number of cycles to response was 2 (range 1–9, mean 4) but three patients needed 8 or 9 cycles. Median response duration was 19 months (4–42). Median survival was 9 months, 23 months for responders, 9 months for marrow responders, and 7 months for non-responders (see Fig. 1a).

Potential factors influencing treatment response

Four patients received hydroxyurea (HU) for more than 1 cycle during their study treatment, due to leukocytosis from start of study treatment. One of them had stable disease, two were classified as having progressive disease, and one patient remained stable until HU was discontinued. The latter patient (no. 5) with CMML-II illustrates that HU may influence the response to epigenetic treatment. He started with a marrow blast count of 15% in March 2007, which was virtually unchanged (18%) prior to cycle 5. Regarding cytogenetics, 100% of mitoses ($n=22$) showed trisomy 8. After 8 cycles, 1 cycle after HU had been stopped, marrow blasts had decreased to 2%. Cells still harbored trisomy 8 [18/18]. At the same time, platelet transfusion dependency had subsided. After 12 cycles, the patient achieved CgCR according to conventional cytogenetics and BM-FISH (46, XY [24/24]) and PR by PB-CD34-FISH and mitochondrial DNA monitoring. Apparently, study treatment only became successful after HU had been discontinued (see Discussion).

We analyzed other factors possibly related to response, as shown in Table 5. There was no significant influence of age, gender, IPSS, primary vs. secondary MDS, WBC, platelets, and karyotype, although four out of seven patients (57%) with chromosome 7 abnormalities responded, including two with a complex karyotype. One further patient with monosomy 7 and complex karyotype achieved a marrow response. Factors negatively influencing response were AML (vs. MDS), marrow blast count, pretreatment, and transfusion dependency. Of the ten patients who had received previous treatment, only one responded, while 8 out of 14 untreated patients (57%) in the entire group and 7 out of 11 (64%) untreated MDS patients responded. Variables influencing survival were AML (vs. MDS) and transfusion dependency.

The influence of VPA serum levels was also evaluated. Median VPA dosage was 25 mg/kg (14–41). The highest

Table 2 Characteristics of responding patients

No.	Age	Diagnosis (WHO, FAB)	IPSS	Karyotype	Pretreatment	Total no. cycles	Response	Cytogenetic response	Cycles to response	Response duration/ months	Dose reductions	Status	Survival
D05 ^a	69	CMML II, CMMML	Int- II	+8	Hydroxyurea	23	CR	CR	9	20	Aza 75%	Dead	26
D03 ^a	73	RAEB II, RAEB	Int- II	Normal	None	18	CRdys	na	2	20	Aza 50%	Dead	23
D04 ^a	72	sAML/MDS RAEB I	Int II	Complex (incl. -7)	None	6	CRdys	PR	1	9	none	Dead	11
D06 ^a	64	CMML II, CMMML	High	del(7q)	None	43	CRdys	CR	2	44	Aza 50%	Alive, ongoing	44
D17 ^a	67	sAML/MDS, RAEB-T	High	del(20q)	None	19	CRdys	CR	9	16	Aza 50%, VPA reduction	Dead	30
F03 ^a	77	AML de-novo	X	+21	Intensive		chemotherapy	10	CRi	CR	8	13	VPA
D14 ^a	77	sAML/MDS, RAEB-T	High	Normal	interruption	Dead	13						
D01 ^a	69	sAML/MDS RAEB	High	Complex (incl -7)	None	4	PR	na	5	29	VPA		
D15 ^a	74	RAEB I, RAEB	Int- II	-7, dup 1 (q)	None	5	PR	discontinuation	19	4	VPA		
F02 ^b	87	AML de-novo	X	Normal	LBH589	3	SD, marrow CR	dead	6	2	VPA interruption	dead	2
F05 ^b	59	sAML/MDS, RAEB-T	High	Complex (incl -7)	Intensive		chemotherapy	3	SD,	marrow CR	none	na	3
D11 ^b	74	t-AML	X	+6,+8	interruption	allo-	PBSCT ->CR, dead	13	na	8	Aza 75%	dead	9
D13 ^b	72	sAML/MDS	X	Normal	None	6	SD, marrow PR (blast count: 34→8%)	na	na	8	none	dead	9

na not applicable, nd not done

^a Responders

^b Marrow remissions only

Table 3 Characteristics of response and treatment

Characteristic	No. (range)
Cytologic CR/PR	7/2
Marrow response CMR/PMR	3/1
Hematologic improvement	
HI-N	6
HI-P	9
HI-E	5
Cytogenetic response (conventional) CR/PR	6/0
FISH response (BM) CR/PR/not done	4/1/1
FISH response (CD34, pB)CR/PR	4/2
Mitochondrial mutations CR/PR	2/1
Median survival	
All patients	9.5 months
Responders vs. marrow response vs. nonresponders	23 vs. 9 vs. 7 months
Median number of cycles	5(2–43)
Median number of cycles to response	2(1–9)
Dose reduction vidaza	9
75%	5
50%	4
Median dosage VPA mg/kg	25 (14–41)
Maximum VPA level (median)/ $\mu\text{g/ml}$	86 (58–122) ^a
Median VPA level (median)/ $\mu\text{g/ml}$	68 (44–97) ^a
Median time to VPA level >70 $\mu\text{g/ml}$	4 weeks ^b
Patients with significant VPA side effects leading to:	13
Dose reduction/Treatment break/cessation VPA	7/5/3

^a not done, $n=1$ ^b never achieved, $n=1$

trough level was 86 $\mu\text{g/ml}$ (58–122). Median VPA trough level over the whole treatment period was 68 $\mu\text{g/ml}$ (44–97), due to necessary dose reductions. A serum level >70 $\mu\text{g/ml}$ was achieved after a median of 4 weeks. The likelihood of response was influenced by median ($p=0.004$) and maximum VPA levels ($p=0.007$). The time to reach a VPA level above 70 $\mu\text{g/ml}$ also provided prognostic information ($p=0.02$). The strongest predictive factor ($p=0.00005$) was the VPA level achieved within the first 2–3 weeks of treatment. The influence of median VPA serum levels on survival is illustrated by Fig. 1c and Table 5. On the other hand, VPA dosage (absolute amount or dose/kg) was not related to response or survival. On multivariate analysis, median VPA level was the only independent prognostic factor for response ($p=0.015$), while MDS vs. AML ($p=0.031$) and transfusion dependency ($p=0.032$) were the only independent parameters influencing survival.

Side effects

Adverse events are summarized in Table 6. Most patients had transient central nervous system (CNS) side effects, necessitating dose reductions or temporary treatment interruption in eight and cessation of therapy in two patients. In one patient, Coombs-negative hemolysis occurred. It remained unclear whether this was related to one of the study drugs. Discontinuation of VPA was without effect, and hemolysis did not respond to steroids. Two patients developed a rash after AZA but were able to continue treatment with concomitant steroids. Myelosuppression occurred in all patients, especially during the initial

Table 4 Response details

Patient	Cytologic response (blast count) (%)	Hematological improvement	Cytogenetic response	Response FISH-BM (%)	Response FISH-CD34 (%)	Mito response (%)
F03	CRi	HI-P	CR	nd	CR	na
D01	PR (5)	HI-N, HI-P, HI-E	nd	nd	nd	nd
D03	CRdys	HI-N, HI-P, HI-E	na	na	na	na
D04	CRdys	HI-N, HI-P, HI-E	CR	CR	CR	na
D05	CR	HI-N, HI-P	CR	CR	PR (1)	PR (5)
D06	CRdys	HI-P	CR	CR	PR (5)	na
D14	CRdys	None	na	na	na	na
D15	PR (5)	HI-P, HI-E	CR	PR (6)	CR	CR
D17	CRdys	HI-N, HI-P, HI-E	CR	CR	CR	CR
F02	SD (CMR)	None	na	na	na	na
F05	SD (CMR)	None	None	nd	none	na
D11	SD (CMR)	None	None	none	none	None
D13	SD (PMR)	None	na	na	na	None

nd not done, na not applicable




Fig. 1 Survival from start of treatment, **a** for the entire cohort, **b** responders vs. nonresponders, landmark analysis at 1 year (at 2 years, 44% of responders are alive compared to 0% of non-responders) **c**) according to VPA serum level $\leq 70 \mu\text{g/ml}$

cycles. Treatment had to be delayed in several patients after the first or second cycle, and dose reductions of AZA were made in seven cases. One patient died of pneumonia after the first cycle.

Discussion

Our study of AZA plus VPA confirms that this combination is feasible and efficacious with an ORR of 37% and a CR rate of 29% in patients with higher-risk MDS and AML. However, since our study was not randomized, the additive

value of VPA cannot be determined. AZA as monotherapy significantly prolongs survival of patients with higher-risk MDS (Fenaux et al. 2009). Still, about 50% of patients do not respond, and response duration is limited despite continued administration. In order to augment the effect of AZA on epigenetic regulation of gene expression, the most promising approach was to combine the demethylating agent with an HDAC inhibitor (Yang et al. 2005).

Most combination studies to date have used VPA, which requires higher concentrations than newer HDAC inhibitors but is relatively selective for class-I-HDACs and leads to degradation of HDAC 2 (Göttlicher et al. 2001). Protocols

Table 5 Comparison of characteristic responders vs. non-responders and influence of different characteristics on response rate and survival

Characteristics	Responders N=8	Nonresponders incl. marrow CR N=16	p response rate	p survival
Median age	72	74	n.s.	n.s. ^a
Sex	78% male	80% male	n.s.	n.s.
Karyotype risk (IPSS)	44% high	53% high	n.s.	n.s.
Normal/abnormal KT	22% normal	33% normal	n.s.	n.s.
Chromosome 7 abnormality	44% Chr 7	27% Chr 7	n.s.	n.s.
IPSS score	50% high	83% high	n.s.	n.s.
MDS vs. AML	89% MDS	33% MDS	0.02	0.0052
Prim vs. sek.	13% sek	33% sek	n.s.	n.s.
Treatment vs. no pretreatment	11% yes	60% yes	0.03	n.s.
Mean marrow blast count	15	40	0.009	n.s. ^b
WBC/ μl	3,000	3,900	n.s.	n.s. ^c
Transfusion dependency	56%	93%	0.047	0.014
Platelet count/ μl	42	76	n.s.	n.s. ^d
Median VPA level (mean)/ $\mu\text{g/ml}$	77	62	0.004	0.03 ^e
Maximum VPA level (mean)/ $\mu\text{g/ml}$	101	80	0.007	0.040 ^f
Time to level $>70 \mu\text{g/ml}$	3	7	0.02	n.s. ^g
VPA level week 2/3	79	46	0.00005	0.0012 ^h
VPA dosage (mg)	1,800	2,066	n.s.	n.s. ⁱ
VPA dosage/kg	23	27	n.s.	n.s. ^j

cutoff (=median):

^a 70 years

^b 20% marrow blasts

^c 3,500/ μl

^d 60.000/ μl

^e 70 $\mu\text{g/ml}$

^f 80 $\mu\text{g/ml}$

^g 5 weeks

^h 50 $\mu\text{g/ml}$

ⁱ 2,000 mg

^j 25 mg/kg

Table 6 Side effects 9 months 0.6%, 77.7% $p=0.0038$

Side effects grades 1/2	Frequency
Local skin reaction	25
Nausea	24
Diarrhea	19
Constipation	17
Leukocytosis	15
Fatigue	14
Hypokaliemia	13
Somnolence	12
Fever	10
Influenza-like symptoms	10
Side effects grade 3/4	Frequency
Neutropenic fever	12
Pneumonia	10
Febrile neutropenia	4
Septicemia	4
Dyspnea	3
Somnolence	3
Catheter-side infection	3
Neutropenia	3
Pancytopenia	3
Confusion	3

vary considerably. All trials except ours added ATRA to the treatment regimen, although Voso et al. (2009) did so only in non- or suboptimal responders. Craddock et al. (2008) added theophylline as a fourth substance. Various AZA dosages have been used. In our trial, we chose a novel dosing regimen for AZA (100 mg/m²/day for 5 days) which avoids the “weekend problem” while providing an overall dose of AZA (500 mg/m²/cycle) that is very similar to the approved regimen (525 mg/m²/cycle). The higher dose intensity of AZA in our schedule may have contributed to the increased myelosuppression compared to our previous experience with AZA monotherapy. However, the combination with VPA, as well as unfavorable patient characteristics, represent other potentially important factors. Two previous studies (Soriano et al. 2007; Raffoux et al. 2010) used a high-dose intermittent schedule for VPA. We chose to apply VPA on a continuous schedule. The doses administered by us and the serum trough levels achieved were higher than in the Italian study (Voso et al. 2009). Despite major differences in these three treatment regimens, the significant influence of VPA serum levels on response rate and survival in our study was the same as that observed by Soriano et al. and Voso et al. Also in line with previous observations, we found no influence of VPA dosage, probably due to pharmacogenomic differences between patients, as

suggested by Voso et al. (2009). As far as typical side effects of HDAC inhibitors, like CNS or gastrointestinal symptoms are concerned, we observed no increased incidence or severity during combination treatment with AZA.

Combination trials have also been performed using decitabine as a demethylating agent. A randomized trial adding VPA intermittently to decitabine in one of the two arms (Issa et al. 2008) yielded response rates (52% vs. 43%) that failed to show a statistically significant difference, probably due to the limited number of patients included ($n=67$).

The overall response rate in our trial was 37%, which is lower than expected from previous trials with AZA monotherapy (Fenaux et al. 2009). However, our series included a high proportion of AML patients, particularly patients pretreated with intensive chemotherapy and/or HDAC inhibitors. Although VPA can be beneficial for patients with AML, the majority of patients included in the abovementioned clinical trials had a relatively low marrow blast count (Fenaux et al. 2009; Sudan et al. 2006). In contrast, the French ATU program included 184 patients with relapsed/refractory AML and generated a response rate of only 13% (Itzykson et al. 2009). In our study, the response rate according to IWG criteria was 57% for all untreated patients (MDS and AML) and 64% for untreated patients with MDS. The other combination trials mentioned above reached comparable response rates. Phase II trials are often difficult to compare, due to varying patient and response characteristics, which are given in Table 7. For example, Voso and coworkers decided to analyze responses only in patients who completed 8 cycles. Some studies, in contrast to our trial, classified marrow responses as response rather than stable disease. Furthermore, IWG criteria, which are currently considered the gold standard for response evaluation, are different for MDS and AML.

A relatively large proportion of our patients achieved CR (29% overall and 55% of previously untreated patients with MDS). Hematological CR was accompanied by complete cytogenetic remission in all evaluable cases. All long-time hematologic improvements were associated with marrow remissions, while not all marrow remissions produced hematologic improvement. An improved CR rate with combination therapy has been described in previous trials (Soriano et al. 2007). Whether improved CR rates were due to the addition of VPA, a slightly increased daily dosage of AZA, or other causes is not entirely clear from these phase II studies. The same is true for time to response, which appeared shorter than with AZA monotherapy in the study reported by Soriano et al. That observation is in accordance with the median time to response in our trial (only 2 cycles). However, even though combination therapy may achieve earlier responses in a proportion of patients, it does not seem to obviate the need for prolonged treatment. While five of our patients responded quite early, after only 1 or

Table 7 Overview of different combination studies with VPA and AZA

Patient	This study	Soriano et al.	Voso et al.	Raffoux et al.	Craddock et al.
Drugs	VPA, 5-Aza	VPA, Aza, ATRA	VPA, Aza, ATRA ^a	VPA, Aza, ATRA	VPA, Aza, ATRA, theophylline
Dosage VPA	Median dosage 25 mg/kg continuously	50–75 mg/kg 7 days	600–1,500 mg continuously	35–50 mg/kg 7 days	ng
Dosage AZA	100 mg/m ² 5 days	75 mg/m ² 7 days	75 mg/m ² 7 days	75 mg/m ² 7 days	75 mg/m ² 5 days
No. of pts.	<i>n</i> =25, evaluable 24	<i>n</i> =53	<i>n</i> =62, 26 evaluable ^b	<i>n</i> =65	<i>n</i> =45
Median age	73 years	69 years	70 years	72 years	66 years
No. AML/MDS	75%/25% ^c	92%/8%	31%/69% ^d	85%/15%	80%/20%
Pretreatment	<i>n</i> =10 (42%)	<i>n</i> =20 (38%)	ng	<i>n</i> =13 (20%)	<i>n</i> =30 (67%)
ORR	37%	27% ^e	31%	26%	33%
CR/PR/Hi/marrow response	7/2/0/4	27/0/0/7	3/5/4/ng	14/3/ng ^f /ng	7/8/ng/ng
Median cycles to response	2 (1–9)	2 (1–3)	5 (2–10) months ^g	ng	2 (1–6)
OS	9 months (23 in responders)	6.5 months (14+ in responders)	14.4 months	12.4 months (19.5 in responders)	ng

ng not given

^a only in non-responders

^b at least 8 cycles to be evaluable

^c 46%/54% according to FAB

^d 100% according to FAB

^e 42% including marrow response

^f 12.5% of non-responders had HI-erythroid

^g Time to CR

2 cycles, three other patients needed 8 or 9 cycles. Like AZA monotherapy, combination therapy with AZA plus VPA requires perseverance, and it is now generally recommended that patients who achieve at least stable disease should be kept on treatment as long as possible.

An interesting finding in our study was that patients can remain transfusion-free with stable peripheral blood counts despite showing a relapse in the bone marrow. We found that increases in medullary blasts or aberrant cells may occur without being followed by clinically relevant progression/relapse. Actually, the two patients with the longest treatment benefit had cytogenetic or even cytomorphological full-blown relapse. These findings illustrate that epigenetic treatment may require changes in our concept of evaluating treatment success. Achieving CR is not paramount, since it is not a prerequisite for obtaining a substantial survival benefit. The AZA-001 trial (Fenaux et al. 2010) made it clear that patients whose best response was hematologic improvement had the same increase in life expectancy as patients who achieved CR or PR.

Even though a number of patients in our trial achieved hematological CR, most of them showed persistent dysplastic features, as it is often the case after chemotherapy in MDS as well. This was even true for patient 17, who was in

complete remission according to conventional cytogenetics, PB-CD34-FISH, BM-FISH, and mitochondrial mutation analysis. This patient relapsed relatively quickly after 16 cycles. This suggests that in some cases, the available monitoring tools may detect only the most advanced, dominant MDS clone without revealing an underlying preleukemic clone that may be responsible for the dysplastic features.

As in previous studies, favorable responses were observed in patients with chromosome 7 abnormalities (Raj et al. 2007). The response rate may not be better than with a normal karyotype or trisomy 8, but this group of patients is clearly doing worse with conventional therapies. In contrast to Mufti et al. (2009), we even observed responses in two of three patients who had chromosome 7 abnormalities as part of a complex karyotype (survival time 11 and 19 months, respectively).

An interesting aspect of our study relates to the possible interaction of hydroxyurea with study medication. In vitro studies suggest that inhibition of ribonucleotide reductase by HU (Choi et al. 2007) interferes with conversion of 5-azacytidine to 5-aza-2'-deoxycytidine, thereby decreasing incorporation of the false nucleotide into DNA and diminishing its hypomethylating effect. Although it has not been proven that DNA hypomethylation is

the only mechanism behind the clinical efficacy of AZA, the clinical course of our patient who achieved remission only after cessation of HU appears to underscore the relevance of HU interference with AZA conversion, thus pointing to the importance of AZA getting incorporated into DNA. It is worth mentioning that three further patients who received HU during study treatment were non-responders.

Our trial was the first to investigate the combination of AZA plus VPA without additional ATRA. Although the response rate was not higher than would be expected with AZA monotherapy, the heterogeneity of the patient population, with a significant proportion of pretreated AML patients, should be taken into consideration. The CR rate was favorable compared with previously published trials, and some of the responses appeared to occur faster than with single-agent AZA. These observations as well as the significant influence of VPA serum levels on response suggest that the HDAC inhibitor contributes to treatment success. However, the true value of HDAC inhibitors in this kind of epigenetic combination therapy can only be determined by randomized trials.

Conflict of interest A. Kuendgen and G. Bug received lecture honoraria and travel support from Celgene; O. Ottmann had an advisory board participation at Celgene; and U. Germing and N. Gattermann received lecture honoraria and research support from Celgene

References

- Bennett JM, Catovsky D, Daniel MT et al (1982) Proposals for the classification of the myelodysplastic syndromes. *Br J Haematol* 51:189–199
- Bhalla KN (2005) Epigenetic and chromatin modifiers as targeted therapy of hematologic malignancies. *J Clin Oncol* 23:3971–3993
- Braulke F, Schanz J, Jung K et al (2010) FISH analysis of circulating CD34+ cells as a new tool for genetic monitoring in MDS: verification of the method and application to 27 MDS patients. *Leuk Res* 34:1296–1301
- Cheson BD, Greenberg PL, Bennett JM et al (2006) Clinical application and proposal for modification of the International Working Group (IWG) response criteria in myelodysplasia. *Blood* 108:419–425
- Choi SH, Byun HM, Kwan JM, Issa JP, Yang AS (2007) Hydroxycarbamide in combination with azacitidine or decitabine is antagonistic on DNA methylation inhibition. *Br J Haematol* 138:616–623
- Craddock C, Goardon N, Griffiths M, Cavenagh JD, Siddique S, Vyas P (2008) 5' Azacitidine in Combination with Valproic Acid Induces Complete Remissions in Patients with Advanced Acute Myeloid Leukaemia but Does Not Eradicate Clonal Leukaemic Stem/Progenitor Cells. *ASH Annual Meeting Abstracts* 112:945
- Fenaux P, Mufti GJ, Hellstrom-Lindberg E et al (2009) Efficacy of azacitidine compared with that of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: a randomised, open-label, phase III study. *Lancet Oncol* 10:223–232
- Fenaux P, Mufti GJ, Hellström-Lindberg E et al (2010) Azacitidine prolongs overall survival compared with conventional care regimens in elderly patients with low bone marrow blast count acute myeloid leukemia. *J Clin Oncol* 28:562–569
- Göttlicher M, Minucci S, Zhu P et al (2001) Valproic acid defines a novel class of HDAC inhibitors inducing differentiation of transformed cells. *EMBO J* 20:6969–6978
- Greenberg P, Cox C, LeBeau MM et al (1997) International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood* 89:2079–2088
- Grimwade D, Walker H, Oliver F et al (1998) The importance of diagnostic cytogenetics on outcome in AML: analysis of 1,612 patients entered into the MRC AML 10 trial. The Medical Research Council Adult and Children's Leukaemia Working Parties. *Blood* 92:2322–2333
- Harris NL, Jaffe ES, Diebold J et al (1999) The World Health Organization classification of neoplastic diseases of the hematopoietic and lymphoid tissues. Report of the Clinical Advisory Committee meeting, Airlie House, Virginia, November, 1997. *Ann Oncol* 10:1419–1432
- Issa JP, Castoro R, Ravandi-Kashani F et al (2008) Randomized phase II study of combined epigenetic therapy: decitabine vs. decitabine and valproic acid in MDS and AML. *ASH Annual Meeting Abstracts* 112:228
- Itzykson R, Thepot S, Recher C et al (2009) Azacytidine in refractory or relapsed AML after intensive chemotherapy (IC): results of the French ATU Program. *ASH Annual Meeting Abstracts* 114:1054
- Kuendgen A, Strupp C, Aivado M et al (2004) Treatment of myelodysplastic syndromes with valproic acid alone or in combination with all-trans retinoic acid. *Blood* 104:1266–1269
- Kuendgen A, Schmid M, Schlenk R et al (2006) The histone deacetylase (HDAC) inhibitor valproic acid as monotherapy or in combination with all-trans retinoic acid in patients with acute myeloid leukemia. *Cancer* 106:112–119
- Kuendgen A, Gattermann N (2007) Valproic acid for the treatment of myeloid malignancies. *Cancer* 110:943–954
- Lubbert M, Schmid M, Ruter B et al (2008) Continued treatment with an outpatient maintenance schedule of decitabine in older AML patients ineligible for induction chemotherapy: results of the 00331 phase II multicenter trial. *ASH Annual Meeting Abstracts* 112:2965
- Maurillo L, Spagnoli A, Genuardi M et al (2008) 5-Azacitidine for the treatment of acute myeloid leukemia: a retrospective, multicenter study of 55 patients. *ASH Annual Meeting Abstracts* 112:1947
- Mufti GJ, Gore SD, Santini V et al (2009) Influence of karyotype on overall survival in patients with higher-risk myelodysplastic syndrome treated with azacitidine or a conventional care regimen. *ASH Annual Meeting Abstracts* 114:1755
- Phiel CJ, Zhang F, Huang EY, Guenther MG, Lazar MA, Klein PS (2001) Histone deacetylase is a direct target of valproic acid, a potent anticonvulsant, mood stabilizer, and teratogen. *J Biol Chem* 276:36734–36741
- Raffoux E, Cras A, Recher C et al (2010) Phase 2 clinical trial of 5-azacitidine, valproic acid, and all-trans retinoic acid in patients with high-risk acute myeloid leukemia or myelodysplastic syndrome. *Oncotarget* 1:34–42
- Raj K, John A, Ho A et al (2007) CDKN2B methylation status and isolated chromosome 7 abnormalities predict responses to treatment with 5-azacytidine. *Leukemia* 21:1937–1944
- Silverman LR, Holland JF, Demakos EP et al (1994) Azacitidine (Aza C) in myelodysplastic syndromes (MDS), CALGB studies 8421 and 8921. *Ann Hematol* 68:A12 (meeting abstract)
- Silverman LR, Demakos EP, Peterson BL et al (2002) Randomized controlled trial of azacitidine in patients with the myelodysplastic

- syndrome: a study of the cancer and leukemia group B. *J Clin Oncol* 20:2429–2440
- Silverman LR, McKenzie DR, Peterson BL et al (2006) Further analysis of trials with azacitidine in patients with myelodysplastic syndrome: studies 8421, 8921, and 9221 by the cancer and leukemia group B. *J Clin Oncol* 24:3895–3903
- Soriano AO, Yang H, Faderl S et al (2007) Safety and clinical activity of the combination of 5-azacytidine, valproic acid, and all-trans retinoic acid in acute myeloid leukemia and myelodysplastic syndrome. *Blood* 110:2302–2308
- Sudan N, Rossetti JM, Shaddock RK, Latsko J, Lech JA, Kaplan RB, Kennedy M, Gryn JF, Faroun Y, Lister J (2006) Treatment of acute myelogenous leukemia with outpatient azacitidine. *Cancer* 107:1839–1843
- Voso MT, Santini V, Finelli C et al (2009) Valproic acid at therapeutic plasma levels may increase 5-azacytidine efficacy in higher risk myelodysplastic syndromes. *Clin Cancer Res* 15:5002–5007
- Wulfert M, Küpper AC, Tappich C et al (2008) Analysis of mitochondrial DNA in 104 patients with myelodysplastic syndromes. *Exp Hematol* 36:577–586
- Yang H, Hoshino K, Sanchez-Gonzalez B, Kantarjian H, Garcia-Manero G (2005) Antileukemia activity of the combination of 5-aza-2'-deoxycytidine with valproic acid. *Leuk Res* 29:739–748